

VHE Gamma Rays from PKS 2155–304

P. M. Chadwick, K. Lyons, T. J. L. McComb, K. J. Orford, J. L. Osborne, S. M. Rayner,
S. E. Shaw, K. E. Turver, and G. J. Wiczorek

Department of Physics, Rochester Building, Science Laboratories, University of Durham,
Durham, DH1 3LE, U.K.

ABSTRACT

The close X-ray selected BL Lac PKS 2155–304 has been observed using the University of Durham Mark 6 very high energy (VHE) gamma ray telescope during 1996 September/October/November and 1997 October/November. VHE gamma rays with energy > 300 GeV were detected from this object with a time-averaged integral flux of $(4.2 \pm 0.7_{stat} \pm 2.0_{sys}) \times 10^{-11} \text{ cm}^{-2} \text{ s}^{-1}$. There is evidence for VHE gamma ray emission during our observations in 1996 September and 1997 October/November, with the strongest emission being detected in 1997 November, when the object was producing the largest flux ever recorded in high-energy X-rays and was detected in > 100 MeV gamma-rays. The VHE and X-ray fluxes show evidence of a correlation.

Subject headings: BL Lacertae objects: individual (PKS 2155–304) — galaxies: active
— gamma rays: observations

1. Introduction

Close X-ray selected BL Lacs (XBLs) are sources of very high energy (VHE) gamma rays at energies above 300 GeV. The BL Lac first detected as a source of VHE gamma rays was Mrk 421 (Punch et al. 1992), following its discovery as a GeV source by the EGRET experiment on the *Compton Gamma Ray Observatory* (Lin et al. 1992). Mrk 421 has been extensively monitored in VHE gamma rays and exhibits complex behaviour, having a low-level quiescent state with flaring on time scales as short as 30 minutes (Gaidos et al. 1996; Zweerink et al. 1997; Takahashi et al. 1998). Mrk 501 is also a source of VHE gamma rays (Quinn et al. 1996), although not detected at GeV energies with EGRET, and also exhibits low-level emission with flaring (Catanese et al. 1997a). In 1997, extremely strong outbursts of TeV emission were detected lasting several months (Deckers et al. 1997, Quinn et al. 1997, Hayashida et al. 1998). The BL Lac 1ES 2344+514 also emits VHE gamma rays, most of the evidence for emission coming from a single outburst (Catanese et al. 1997b, Catanese et al. 1998). All of these objects are close ($z \sim 0.03$), X-ray selected BL Lacs. There have been no reported detections of VHE gamma rays from radio-selected BL Lacs.

(RBLs — see e.g. Kerrick et al. 1995, Roberts et al 1998) although this category of object is frequently detected at GeV energies by the EGRET experiment.

Stecker et al. 1996 have interpreted the gamma ray results in the GeV – TeV range and propose a model in which RBLs will be GeV gamma ray sources and XBLs will be TeV sources. They associate the emission of TeV γ -rays from XBLs with the presence of relativistic electrons with energies higher than those in RBLs. They then go on to show that a simple synchrotron self-Compton model can explain the differences observed between RBLs and XBLs provided the attenuation of the VHE gamma ray flux by pair production with the intergalactic infrared background is taken into account (Stecker et al. 1992; Stecker & de Jager 1997a). On the basis of this model, PKS 2155–304, despite having a redshift of 0.117, is predicted to be a strong TeV gamma-ray source.

The BL Lac PKS 2155–304 was discovered as an X-ray source during observations made with the *HEAO-1* satellite (Schwartz et al. 1979; Griffiths et al. 1979) at a position where the *Ariel V* satellite had previously detected confused emission (Cooke et al. 1978). In many ways, PKS 2155–304 may be regarded as the archetypal X-ray selected BL Lac object; like most BL Lac objects it is associated with a compact, flat spectrum radio source and has an almost featureless continuum which extends from radio to X-ray energies. It is the brightest known BL Lac at UV wavelengths, and the object’s maximum power is emitted between the UV and the soft X-ray range (Wandel & Urry 1991). PKS 2155–304 has a history of rapid, strong broadband variability and has been the subject of several multiwavelength monitoring campaigns (see e.g. Brinkmann et al 1994; Courvoisier et al. 1995; Pesce et al. 1997). In 1997 November, contemporaneous with some of the observations reported here, X-ray emission was detected with the *Beppo-SAX* satellite (Chiappetti & Torroni 1997) with a flux equal to the strongest previous outburst.

The EGRET experiment on board the *Compton Gamma Ray Observatory* was used to detect 30 MeV – 10 GeV gamma ray emission from PKS 2155–304 during 1994 November 15 – 29 (Vestrand et al. 1995). These observations indicated a very hard spectrum, with an integral power-law spectral index of 1.71 ± 0.24 , and this, combined with its proximity, make it an excellent candidate TeV source. However, a recent detailed analysis of the shape of the X-ray spectrum suggests that PKS 2155–304 may not be an important TeV emitter, producing photons with a maximum energy of ~ 800 GeV, although a very high degree of beaming may allow the production of photons with higher energies (Giommi et al. 1998). Indeed, the absence of any signature of γ -ray absorption due to pair production at energies up to 8 GeV in observations of PKS 2155–304 is consistent with the suggestion that it contains a relativistic jet oriented nearly along our line of sight (Vestrand et al. 1995).

2. Observations

2.1. The Mark 6 Telescope

The University of Durham Mark 6 atmospheric Čerenkov telescope has been in operation at Narrabri, NSW, Australia since July 1995. The telescope is described in detail elsewhere (Armstrong et al. 1997). It uses the well-established imaging technique to separate VHE gamma rays from the cosmic ray background, combined with a robust noise-free trigger based on the signals from three parabolic flux detectors of 7 m diameter and aperture $f/1.0$ mounted on a single alt-azimuth platform. A 109-element imaging camera (0.25° pixel size) is mounted at the focus of the central flux detector, with low resolution cameras each consisting of 19 pixels (0.5° pixel size) mounted at the focus of the outer (left and right) flux collectors. The telescope is triggered by demanding a simultaneous temporal (10 ns gate) and spatial (0.5° aperture) coincidence of the Čerenkov light detected in the three cameras. This triggering system enables the telescope to detect low energy gamma rays with high immunity from triggering by local muons. Initial work suggests that our system has a detection probability for gamma rays of about 1% at 100 GeV with the probability of detection rising slowly to about 40% at 250 GeV.

The Mark 6 telescope has been designed to provide stable operation which allows the observation of weak DC sources. The multiple-dish triggering system is stable against variations in performance due to accidental coincidences. All detector packages are thermally stabilised. The atmospheric clarity is continuously monitored both using a far infra-red radiometer and an axial optical CCD camera which enables the position and visual magnitude of guide stars to be monitored. The gain and noise performance of the PMTs, digitizer pedestals and associated electronics are continuously monitored by:

1. triggering the telescope at random times using a nitrogen laser / plastic scintillator / optical fibre light guide / opal diffuser system to simulate Čerenkov flashes and so enable flat-fielding, and
2. producing false triggers at random times to measure samples of the background noise.

PKS 2155–304 was observed in 1996 September/October/November and 1997 October/November under moonless, clear skies. The 1997 November observations were made as part of a multiwavelength campaign co-ordinated by T. Vestrand. An observing log is shown in Table 1. Data were taken in 15-minute segments. Off-source observations were taken by alternately observing regions of sky which differ by ± 15 minutes in right ascension from the position of PKS 2155–304 to ensure that the on and off segments possess identical azimuth and zenith profiles. This off-source – on-source – on-source – off-source observing pattern is routinely used to eliminate any first order changes in count rate due to any residual secular changes in atmospheric clarity, temperature etc. The gross counting rate of the telescope was typically ~ 800 per minute near culmination (zenith angle 5°) reducing to ~ 150 per minute at zenith angles of 60° . Most of the observations at large zenith angles were made in 1997 November, in order to maximize exposure during the

Date	No. of scans	Date	No. of scans
	ON source		ON source
1996 September 3	2	1997 October 20	3
1996 September 5	7	1997 October 22	7
1996 September 6	5	1997 October 23	7
1996 September 7	6	1997 October 25	3
1996 September 8	4	1997 October 29	4
1996 September 10	8	1997 October 30	2
1996 September 12	3	1997 November 18	2
1996 September 14	7	1997 November 19	4
1996 September 15	7	1997 November 20	5
1996 September 17	5	1997 November 22	2
1996 September 18	3	1997 November 23	4
1996 October 9	4	1997 November 24	4
1996 October 10	8	1997 November 25	2
1996 October 12	5		
1996 October 13	4		
1996 October 14	6		
1996 November 2	3		
1996 November 7	6		
1996 November 10	4		
1996 November 11	5		
1996 November 12	5		

Table 1: Observing log for our observations of PKS 2155–304 during 1996 and 1997.

multiwavelength campaign. Detections of Mrk 501 during its 1997 flaring activity using the Mark 6 telescope at zenith angles of $> 70^\circ$ have demonstrated the usefulness of TeV observations using the atmospheric Čerenkov technique at very large zenith angles (Chadwick et al. 1998c).

Data were accepted for analysis only if:

1. the sky was clear and stable, and
2. the gross counting rates in each on-off pair were consistent at the 2.5σ level.

When these criteria are met, we have a total of 41 hours of on-source observations with an equal quantity of off-source data.

3. Results

Routine reduction and analysis of accepted data comprises the following steps:

1. calibration of the gains and pedestals of all 147 PMTs and digitizer electronics within a 15 minute segment, using the embedded laser and false coincidence events,
2. software padding of the data to equalize the effects of on- and off-source photomultiplier noise on data selection (Cawley et al. 1993; Fegan 1997),
3. identification of the location of the source in the camera’s field of view for each event, using the axial CCD camera,
4. a calculation of the spatial moments of each shower image relative to the source position, and
5. rejection of events containing an image which would be unlikely to be produced by gamma rays.

Events considered suitable for analysis are those which are confined within the sensitive area of the camera (i.e. within 1.1° of the centre of the camera) and which contain sufficient information for reliable image analysis, i.e. which have $SIZE > 500$ digital counts, where 3 digital counts ~ 1 photoelectron, and 200 digital counts are produced by a 125 GeV gamma ray.

Monte Carlo simulations indicate that the image shape of the Čerenkov light from a γ -ray shower can be approximated by an ellipse, the major axis of which is oriented towards the source position, whereas a cosmic ray (hadronic) shower produces a broader, more irregularly shaped image (Hillas 1985). The image can be parameterized using techniques developed by the Whipple group which describe both the shape and the orientation of the image. In addition a measure of the fluctuations between the centroids of the samples recorded by the left and right flux collectors of the Mark 6 telescope provides a further discriminant, D_{dist} (Chadwick et al. 1998a). Gamma rays are identified on the basis of image shape and left/right fluctuation, and then plotting the number of events as a function of the pointing parameter $ALPHA$; γ -ray events from a point source will appear as an excess of events at small values of $ALPHA$.

The selection criteria applied to these data are summarized in Table 2. They constitute a standard set of criteria developed to include allowance for the variation of parameters with event size and are routinely applied to data from all objects recorded at zenith angles less than 45° . Small changes in the values for $WIDTH$, $CONCENTRATION$ and D_{dist} are made for data recorded at zenith angles greater than 45° . These selection criteria differ little from those used in our earlier reports of the detection of VHE gamma-rays from PSR B1706–44 (Chadwick et al. 1998b) and Cen X-3 (Chadwick et al. 1998a); a decrease in the weight given to the signal recorded from the outer (guard ring) detectors in the camera has produced an $ALPHA$ -plot for the background with an increasing population of large $ALPHA$ events.

Parameter	Ranges	Ranges	Ranges	Ranges	Ranges
<i>SIZE</i> (d.c.)	500 – 800	800 – 1200	1200 – 1500	1500 – 2000	2000 – 10000
<i>DISTANCE</i>	0.35° – 0.85°	0.35° – 0.85°	0.35° – 0.85°	0.35° – 0.85°	0.35° – 0.85°
<i>ECCENTRICITY</i>	0.35 – 0.85	0.35 – 0.85	0.35 – 0.85	0.35 – 0.85	0.35 – 0.85
<i>WIDTH</i>	< 0.10°	< 0.14°	< 0.19°	< 0.32°	< 0.32°
<i>CONCENTRATION</i>	< 0.80	< 0.70	< 0.70	< 0.35	< 0.25
D_{dist}	< 0.18°	< 0.18°	< 0.12°	< 0.12°	< 0.10°

Table 2: The image parameter selections applied to the PKS 2155–304 data recorded at zenith angles less than 45°. For data recorded at zenith angles greater than 45° small modifications are made to the *WIDTH*, *CONCENTRATION* and D_{dist} criteria.

The number of events remaining ON and OFF source after the application of the selections described above are summarized in Table 3. The *ALPHA* distribution for the whole dataset, after subtracting the results for OFF from ON observations, is shown in Figure 1, together with the *ALPHA* distributions for the ON and OFF source data individually. No normalization of ON and OFF data rates has been applied. There is an excess of events at small *ALPHA*, the expected γ -ray domain, and imposing a selection of *ALPHA* < 22.5° yields a gamma ray detection significant at the 6.8 σ level for the total dataset.

	On	Off	Difference	Significance
Number of events	1021083	1023280	–2197	–1.5 σ
Number of size and distance selected events	600856	598733	2123	1.9 σ
Number of shape selected events	37125	36151	974	3.6 σ
Number of shape and <i>ALPHA</i> selected events	6099	5370	729	6.8 σ

Table 3: The results of various event selections for the PKS 2155–304 data. Data from observations at all zenith angles have been combined.

In addition, a false source analysis has been performed by re-analysing the ON and OFF-source data using a matrix of assumed source positions in celestial co-ordinates. For each trial source position in the matrix, *DISTANCE* and *ALPHA* are recalculated for each event, and a




Fig. 1.— (a) The *ALPHA* distributions ON and OFF source for PKS 2155–304. The grey line refers to ON source data. (b) The difference in the *ALPHA* distributions for ON and OFF source events.

selection of events is made with values of *DISTANCE* as above and *ALPHA* $< 22.5^\circ$. The result is shown in Figure 2; the excess of gamma ray candidates maximises when the source is assumed to be at the centre of the plot. The central point in the matrix is the position of PKS 2155–304; it is not the geometrical centre of the camera, which is typically $0.1^\circ - 0.2^\circ$ from the source position.

3.1. Observed flux

544 excess events identified as γ -rays were detected in 32.5 hours of on-source observation at zenith angles less than 45° . The current selection procedure for data recorded at zenith angles $< 45^\circ$ is estimated to retain $\sim 20\%$ of the original gamma ray events. The collecting area has been estimated from Monte Carlo simulations and is $\sim 5.5 \times 10^8 \text{ cm}^2$ for these observations. Using this estimate, the γ -ray flux incident on the earth’s atmosphere was $(4.2 \pm 0.75_{\text{stat}} \pm 2.0_{\text{sys}}) \times 10^{-11} \text{ cm}^{-2} \text{ s}^{-1}$. The average gamma ray threshold for this observation, defined as the energy at which the trigger probability is $\sim 50\%$, was 300 GeV.

We also note that the high energy (large size) bin for our observations at zenith angles greater




Fig. 2.— The significance of the number of excess events as a function of assumed source position for PKS 2155–304. The grey scale is such that black corresponds to a detection probability of $> 6\sigma$. Contours are at 0.7σ intervals.

than 45° yields a detection significant at the 2σ level. We estimate that this sample corresponds to a minimum γ -ray energy of about 3 TeV and so there is evidence for detection of photons with energies > 3 TeV from PKS 2155–304.

3.2. Time variability

A characteristic of previous observations of VHE gamma rays from XBLs has been the strong time variability. We have investigated our data for variations in the TeV flux from PKS 2155–304 on timescales of months and days.

We show in Figure 3 the variation in the detected γ -ray flux averaged over the observations made in the observing periods in 1996 September, October and November and in 1997 October and November. The strength of the γ -ray emission is defined as the number of γ -ray candidates (shape




Fig. 3.— The variation of the VHE gamma ray flux from PKS 2155–304 averaged over the observing periods in 1996 September, October and November and in 1997 October and November.

and orientation selected events) to the number of background protons during the same observation. This makes some allowance for variations in the sensitivity of our telescope as the zenith angle changes. Using the test for constancy of emission used by the EGRET group (McLaughlan et al. 1996), the data suggest that the VHE γ -ray emission is time variable. The largest excursion from the average flux occurred during observations in 1997 November.

The variation of the gamma-ray emission during observations made in individual nights is shown in Figure 4. We have no evidence for strong gamma-ray flaring on a time scale of days during these observations. However, the data suggest that the VHE gamma-ray emission is variable on a timescale of days, using the test of McLaughlan et al. 1996.

4. Discussion

PKS 2155–304 is the fourth X-ray emitting BL Lac to be established as a VHE emitter and, thus far, the furthest from earth with a redshift of 0.117. The emission shows the features of time variability demonstrated in the other BL Lacs detected at VHE energies.

In Figure 5 we show the spectral energy distribution (SED) from PKS 2155–304 from radio to VHE gamma ray energies, including the present results. The present VHE result is consistent with the SEDs for the other VHE-emitting blazars. The VHE behaviour is in general agreement with the predictions of Ghisellini et al. 1998 and, as for the other three established VHE blazars, indicates the success of the unified model of AGNs in explaining the gross features of VHE emission. However, the present VHE data are unable to distinguish between the predictions for the external Compton or the synchrotron self-Compton model, as is the case for TeV observations of the other




Fig. 4.— The variation of VHE gamma ray flux from PKS 2155–304 from each observation made in (a) 1996 September, October and November and (b) 1997 October and November.

VHE blazars.

During our observations of PKS 2155–304 measurements of its 2 – 10 keV X-ray emission were available from the ASM on *RXTE*.¹ In Figure 6(a) we show the correlation between the average VHE emission during each of the five months when data were available and the corresponding average X-ray rate from the *RXTE* quick look analysis. The correlation coefficient for the data is 0.66. A similar plot for the daily averaged TeV and X-ray measurements are shown in Figure 6(b). In this case, the correlation coefficient is 0.22.

The data in Figure 6 show evidence for a correlation between the strengths of VHE gamma ray emission and X-ray emission from PKS 2155–304 on timescales of both days and months. In particular, the observing periods during which we failed to detect significant VHE gamma-ray emission (1996 October and November) coincide with a period of low X-ray emission according to the *RXTE* ASM measurements. We also note that Giommi et al. 1998 based their conclusions that PKS 2155–304 should not emit VHE gamma rays from an X-ray observation taken with *Beppo-SAX* shortly after our 1996 November (null) observation. The general correlation between X-ray and VHE emission in PKS 2155–304 is in agreement with multiwavelength observations of Mrk 421 and Mrk 501 (reviewed by e.g. Ulrich, Maraschi & Urry 1997). These show such correlated behaviour, indicating that the same population of electrons produces the X-ray and TeV emission; this is the basis for many models for VHE production in AGNs.

As has been pointed out by Catanese et al. 1997a and Dermer 1998, a detailed study of the temporal correlation between the X-rays and TeV gamma-rays during flares can provide a test for beaming. However, none of the flaring activity observed in our observations of PKS 2155–304 is at

¹Available on the web at <http://space.mit.edu/XTE/asmlc/pks2155-304.html>.




Fig. 5.— The spectral energy distribution of PKS 2155–304. The VHE point is from the present work.

a strength where meaningful limits can be placed on the contribution of beaming.

The strongest VHE emission from PKS 2155–304 occurred in 1997 November (during the multiwavelength campaign). We note that the strongest X-ray emission ever observed from this object was detected by *Beppo-SAX* during observations in 1997 November 22 – 24 (Chiappetti & Torroni 1997). EGRET also detected GeV gamma-rays, again at a flux considerably higher than previous detections, during an observation from 1997 November 11 – 17, just before the onset of our November observation (Sreekumar & Vestrand 1997). The results of the 1997 November multiwavelength campaign will be reported elsewhere.

Our observations indicate that PKS 2155–304 emits VHE gamma rays at energies above about 300 GeV and that the energy spectrum extends, with significance at the 2σ level, beyond 3 TeV. The estimated average integral flux above a threshold of 300 GeV is $(4.2 \pm 0.7_{stat} \pm 2.0_{sys}) \times 10^{-11} \text{ cm}^{-2} \text{ s}^{-1}$.

Observations of VHE gamma-rays from AGNs can be used to probe the inter-galactic infra-red radiation field (Stecker et al. 1992). The most recent version of this model (Stecker & de Jager 1997b, Stecker & de Jager 1998) uses estimates of the infra-red radiation field due to Malkan & Stecker 1988 to predict the optical depth for TeV sources as a function of distance. They predict




Fig. 6.— The correlation between VHE emission from PKS 2155–304 and the X-ray flux measured by the *RXTE* satellite. Fig 6(a) shows the correlation when averaged over an observing period (~ 10 days); the correlation coefficient is 0.66. Fig 6(b) shows the day-to-day correlations; the correlation coefficient is 0.22.

an optical depth of 1 – 2 for an object emitting 3 TeV gamma-rays at $z = 0.12$. Our detection of > 3 TeV gamma-rays from PKS 2155–304 is thus not in conflict with the predictions of Stecker & de Jager 1998. However, a more sensitive test must await a determination of the VHE gamma-ray spectrum from our data.

We are grateful to the UK Particle Physics and Astronomy Research Council for support of the project and the University of Sydney for the lease of the Narrabri site. The Mark 6 telescope was designed and constructed with the assistance of the staff of the Physics Department, University of Durham. GJW acknowledges the receipt of a Tempus studentship. The efforts of Mrs. S.E. Hilton and Mr. K. Tindale are acknowledged with gratitude. This paper uses quick look results provided by the ASM/*RXTE* team and uses the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, Caltech, under contract with the National Aeronautics and Space Administration.

REFERENCES

- Armstrong, P., et al. 1997, *Exp. Astron.*, in press
- Brinkmann, W., et al. 1994, *A&A*, 288, 433
- Catanese, M., et al. 1997a, *ApJ*, 487, L143
- Catanese, M. A., et al. 1997b, in *Proc. 25th Int. Cosmic Ray Conf.*, Durban, ed. M. S. Potgieter,

- B. C. Raubenheimer, & D. J. van der Walt (Potchefstroom: Potchefstroom University for CHE), Vol. 3, 277
- Catanese, M. A., et al. 1998, *ApJ*, 501, 616
- Cawley, M. F., et al. 1993, in *Towards a Major Atmospheric Cerenkov Detector II*, ed. R. C. Lamb, 176
- Chadwick, P. M., et al. 1998a, *ApJ*, 503, 391
- Chadwick, P. M., et al. 1998b, *Astroparticle Phys.*, 9, 131
- Chadwick, P. M., et al. 1998c, in preparation
- Chiappetti, L., & Torroni, V. 1997, *IAU Circular* 6776
- Cooke, B. A., et al. 1978, *MNRAS*, 182, 489
- Courvoisier, T. J.-L., et al. 1995, *ApJ*, 438, 108
- Deckers, T., et al. 1997, in *Proc. 25th Int. Cosmic Ray Conf.*, Durban, ed. M. S. Potgieter, B. C. Raubenheimer, & D. J. van der Walt (Potchefstroom: Potchefstroom University for CHE), Vol. 3, 245
- Dermer, C. D. 1998, *ApJ*, 501, L157
- Fegan, D. J. 1997, *J. Phys. G. Nucl. Part. Phys.*, 23, 1013
- Ghisellini, G., et al. 1998, *astro-ph/9807317*
- Gaidos, J. A., et al. 1996, *Nature*, 383, 319
- Giommi, P., Ansari, S. G., & Micol, A. 1995, *A&AS*, 109, 267
- Giommi, P., et al. 1998, *A&A*, 333, L5
- Griffiths, R. E., et al. 1979, *ApJ*, 234, 810
- Hayashida, N., et al. 1998, *ApJ*, 504, L71
- Hillas, A. M. 1985, in *Proc. 19th Int. Cosmic Ray Conf.*, La Jolla, ed. F. C. Jones, J. Adams, & G. M. Mason (Washington: NASA), Vol. 3, 445
- Kerrick, A. D., et al. 1995, *ApJ*, 452, 588
- Lamer, G., Brunner, H., & Staubert, R. 1996, *A&A*, 311, 384
- Lin, Y. C., et al. 1992, *ApJ*, 401, L61
- Malkan, M. A., & Stecker, F. W. 1998, *ApJ*, 496, 13

- McLaughlan, M. A., et al. 1996, *ApJ*, 473, 763
- Quinn, J., et al. 1996, *ApJ*, 456, L83
- Quinn, J., et al. 1997, in *Proc. 25th Int. Cosmic Ray Conf.*, Durban, ed. M. S. Potgieter, B. C. Raubenheimer, & D. J. van der Walt (Potchefstroom: Potchefstroom University for CHE), Vol. 3, 249
- Pesce, J.E., et al 1997, *ApJ*, 486, 770
- Punch, M., et al. 1992, *Nature*, 358, 477
- Roberts, M. D., et al. 1998, *A&A*, 337, 25
- Sreekumar, P., & Vestrand, T. 1997, *IAU Circular* 6774
- Stecker, F. W., & de Jager, O. C. 1997a, *ApJ*, 476, 712
- Stecker, F. W., & de Jager, O. C. 1997b, in *Towards a Major Atmospheric Cerenkov Detector - V*, ed. O. C. de Jager (Potchefstroom: Potchefstroom University for CHE), 39
- Stecker, F. W., & de Jager, O. C. 1998, *A&A*, 334, L85
- Stecker, F. W., de Jager, O. C., & Salamon, M. H. 1992, *ApJ*, 390, L49
- Stecker, F. W., de Jager, O. C., & Salamon, M. H. 1996, *ApJ*, 473, L75
- Stickel, M., et al. 1991, *ApJ*, 374, 431
- Schwartz, D. A., et al. 1979, *ApJ*, 229, L53
- Takahashi, T., et al., 1998, *IAU circular* 6888
- Ulrich, M.-H., Marcaschi, L., & Urry, C. M. 1997, *ARA&A*, 35, 445
- Vestrand, W. T., Stacy, J. G., & Sreekumar, P. 1995, *ApJ*, 454, L93
- Wandel, A., & Urry, C. M. 1991, *ApJ*, 367, 78
- Zweerink, J. A., et al. 1997, *ApJ*, 490, L141